

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number:

0 188 256 B1

12

EUROPEAN PATENT SPECIFICATION

- 49 Date of publication of patent specification: **21.08.91** 51 Int. Cl.⁵: **C07C 327/00, C07C 249/00, C07D 207/46, A61K 49/02, C07C 261/00, G01N 33/534, G01N 33/58**
- 21 Application number: **86100360.6**
- 22 Date of filing: **13.01.86**

54 Metal radionuclide labeled proteins for diagnosis and therapy.

- 30 Priority: **14.01.85 US 692000**
- 43 Date of publication of application: **23.07.86 Bulletin 86/30**
- 45 Publication of the grant of the patent: **21.08.91 Bulletin 91/34**
- 64 Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE
- 56 References cited:
EP-A- 0 135 160

JOURNAL OF NUCLEAR MEDICINE, vol. 25,
no. 2, February 1984, pages 223-229, New
York, US; R.F. SCHNEIDER et al.:
N,N'-bis(S-benzoylmercaptoacetamido)ethyl-
enediamine and propylenediamine ligands
as renal function imaging agents. I. Alter-
nate synthetic methods"

- 73 Proprietor: **NeoRx**
410 West Harrison Street
Seattle Washington 98119(US)
- 72 Inventor: **Fritzberg, Alan R.**
16703, 74th Place West
Edmonds, WA 98020(US)
- 74 Representative: **Glawe, Delfs, Moll & Partner**
Patentanwälte
Postfach 26 01 62 Liebherrstrasse 20
W-8000 München 26(DE)

EP 0 188 256 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

EP 0 188 256 B1

Description

BACKGROUND OF THE INVENTION

5 Field of the Invention

Radiolabeled compounds are important tools in medical diagnosis and treatment. Such compounds are employed in a variety of techniques including the diagnosis of deep venous thrombi, the study of lymph node pathology, and the detection, staging and treatment of neoplasms. A number of these compounds employ metal radionuclides such as Technetium-99m. When employing radionuclides for in vivo administration it is desirable that the radionuclide localize in a target organ or cancer site. Therefore, radionuclides are usually formulated to provide preferential binding to or absorption by the particular organ or tissue. There is considerable interest in being able to accurately direct a radionuclide to a preselected site to reduce background radiation directed to surrounding or distant tissue, reduce the dosage, minimize background for in vivo imaging, and minimize undesirable side effects. Toward this end, methods involving specific ligands or receptors to which the radionuclide may be conjugated are of interest.

Description of the Prior Art

References of interest include Khaw et al., J. Nucl. Med. (1982) 23:1011; Rhodes, B.A., Sem. Nucl. Med. (1974) 4:281; Davidson et al., Inorg. Chem. (1981) 20:1629; and Byrne and Tolman, J. Nucl. Med. (1983) 24:P126. See particularly Fritzberg et al., J. Nucl. Med. (1982) 23:592; Fritzberg et al., ibid. (1981) 22:258; and Fritzberg et al., ibid. (1982) 23:P17 for descriptions of mercaptoacetyl derivatives of ethylene diamine carboxylic acid derivatives. See also U.S. Patent Nos. 4,434,151, 4,444,690, and 4,472,509.

25 SUMMARY OF THE INVENTION


Metal radionuclide labeled proteins are provided for the diagnosis and treatment of a variety of pathologic conditions. Specifically, chelated radionuclide protein conjugates are employed for the diagnosis of conditions including lymph node pathology and deep venous thrombi and the detection and staging of neoplasms. Also, chelated radionuclides as protein conjugates are employed for radiotherapy of tumors.

DESCRIPTION OF THE SPECIFIC EMBODIMENTS

Improved methods and compositions are provided related to metal radionuclide chelates, their active esters for conjugating to proteins, and the resulting peptide conjugates, as well as the use of the conjugates in radioimaging and radiotherapy.

The metal chelating compounds will be dithio, diamino- or diamidocarboxylic acids or amines or derivatives thereof, e.g., a N,N'-bis-mercaptoacetyl ω , (ω -x)-diamino carboxylic acid, (x is 1 or 2) esters capable of forming an amide bond in an aqueous medium with a polypeptide, and intermediates to the chelate. The chelating compounds are referred to as N_2S_2 ligands or chelates.

The compounds of this invention have the following formula:



55 wherein:

M is a metal or metal oxide radionuclide ion capable of being chelated;

one of Z¹, Z², Z³, Z⁴, is R'CW'(HNV)_n, Y', and the others are H₂ or =O;

R' is an aliphatic divalent radical of from 1 to 6 carbon atoms and 0 to 2 heteroatoms;

EP 0 188 256 B1

Y' is the leaving group of an active ester, $-\text{NH}_2$, $-\text{NHNH}_2$, or a polypeptide of at least about 1000 molecular weight;

V' is of the formula $\text{R}'\text{CW}'$;

W' is $=\text{NH}$ or $=\text{O}$, with the proviso that the W' bonded to the carbon atom bonded to Y' is H_2 when Y' is $-\text{NH}_2$;

X' is a bond, methylene or CHZ^4 ;

(A)'s are the same or different and are hydrogen or lower alkyl of from 1 to 6 carbon atoms; and
n' is 0 or 1.

Furthermore, R' is an aliphatic divalent radical of from 1 to 6, usually from 1 to 3 carbon atoms, having from 0 to 1 site of aliphatic unsaturation and 0 to 2 heteroatoms, usually straight chain and preferably methylene or polymethylene of from 2 to 3 carbon atoms;

V' is $\text{R}'\text{CW}'$, where the two $\text{R}'\text{CW}'$ s may be the same or different, usually being of from 1 to 7, more usually of from 1 to 6 carbon atoms, preferably of from 2 to 3 carbon atoms; and


Y' is an ester of an hydroxylic compound, where the ester is capable of forming an amide bond with a polypeptide in an aqueous medium without denaturation of the polypeptide; $-\text{NH}_2$; $-\text{NHNH}_2$; an amino acid, or a polypeptide usually of at least about 1000 molecular weight, more usually at least about 2000 molecular weight, generally less than about 1.6 MDal, more usually less than about 800 KDal. Of particular interest are immunoglobulins or specific binding fragments thereof.

A variety of metals may be employed as the radionuclide M. These metals include copper, e.g., ^{67}Cu , and ^{64}Cu , technetium, e.g. $^{99\text{m}}\text{Tc}$; rhenium, e.g. ^{186}Re and ^{188}Re .

The esters are those esters which are known to provide for the reaction with a polypeptide in aqueous medium. One or another of the esters may be preferred, depending upon the particular radionuclide, the protein, and the conditions for conjugation. Common esters which find use are the o- and p-nitrophenyl, 2-chloro-4-nitrophenyl, cyanomethyl, 2-mercaptopyridyl, hydroxybenztriazole, N-hydroxysuccinimide, trichlorophenyl, tetrafluorophenyl, o-nitro-p-sulfophenyl, N-hydroxyphthalimide and the like. For the most part, the esters will be of activated phenols, particularly nitro-activated phenols and cyclic compounds based on hydroxylamine. As other hydroxylic compounds become available, these also may find use in this invention.

The polypeptide compounds may be varied widely, depending upon the nature of the use of the radionuclide. Thus, the compounds may be ligands or receptors. Ligands may include such a variety of compounds as hormones, lymphokines, growth factors, substrates, particularly compounds binding to surface membrane receptors, where the complex may remain bound to the surface or become endocytosed. Among receptors are surface membrane receptors, antibodies, enzymes, naturally occurring receptors, lectins, and the like. Of particular interest are immunoglobulins or their equivalent, which may involve Fab fragments, $\text{F}(\text{ab}')_2$, F., T-cell receptors, etc.

The invention is furthermore directed to compounds of the following formula:



wherein:

one of Z^1 , Z^2 , Z^3 or Z^4 is $\text{RCW}(\text{HNV})_n\text{Y}$, and the others are H_2 or $=\text{O}$;

R is a divalent organic radical of from 1 to 6 carbon atoms and 0 to 2 heteroatoms;

W is $=\text{NH}$ or $=\text{O}$, with the proviso that the W bonded to the carbon atom bonded to Y is H_2 when Y is $-\text{NH}_2$;

V is the same or different from RCW and comes within the definitions of RCW;

n is 0 or 1;

T is a removable protective group, hydrogen, or an alkali metal ion;

Y is an organic oxy compound forming an active ester, $-\text{NH}_2$, $-\text{NHNH}_2$, or a polypeptide of at least two amino acids;

X is a bond, methylene, or CHZ^4 ; and

EP 0 188 256 B1

butyl mercaptan or the like. The ester may now be hydrolyzed to the acid and the metal chelate formed or the thioether reacted with an activated sulfonyl chloride followed by treatment with thioglycolate. Alternatively α -alkylthio substituted acyl compounds may be used with carbodiimide for acylation, followed by cleavage of the thioether with formation of disulfide and reduction of the disulfide to mercapto, as described above.

5 An alternative approach, employed for the 4,5-diaminopentanoate employs the readily available glutamate. After forming the 5- carboxy ester, the amino group is protected and the acid group (1-carboxyl) preferentially reduced to the alcohol. The alcohol is transformed into an active cleaving group, e.g. halide or pseudohalide, followed by displacement with a nitrogen anion, e.g. azide, which serves as an intermediate to the amino group. After catalytic reduction of the amino intermediate to amino and hydrolysis of the ester, 10 the amino groups are acylated with S-protected α -mercaptoacyl groups. The protective groups may be removed, exchanged or otherwise modified, e.g. by introduction of water solubilizing groups.

Various synthetic procedures may be employed for preparing the different N_2S_2 chelate rings. Carboxamides may be formed and reduced using aluminum or borohydrides to form the amine. Amines 15 may be alkylated with aliphatic halides. Ethylene or propylene diamines or carboxyalkylalkylene diamines may be used to link thioglycolic acids. Other synthetic procedures may also be employed depending on the N_2S_2 ligand of interest.

The imidate may be employed by preparing the nitrile of the amino protected α , (ω -x)-diaminoalkyl halide or pseudohalide by displacement with nitrile, mercaptoacylation of the deprotected amino groups as 20 described previously and imidoester formation by conventional techniques, e.g. acidic (HCl) anhydride in alcohol.

The S-protective groups may be varied widely, being acyl groups, thio groups or other compound which provides protection of the thio group during the subsequent manipulations and can be readily removed without deleterious effect on the peptide conjugate.

Illustrative groups include benzoyl, acetyl, m- or p-phthaloyl, thioglycolic, o-carboxythiophenol, ethylthiocarbonate, β -mercaptopropionic, tetrahydropyranyl, sulfonate, etc. Alternatively cyclic di- or polysulfides 25 may be formed. Disulfides may be prepared using sulfinyl halides, dinitrothiophenoxide substituted mercaptans, with mild oxidation in the presence of excess of the protective group, etc.

The protective groups may be removed in a variety of ways. Thioesters may be hydrolyzed using aqueous ammonia, sodium alkoxide in alcohol, or any conventional technique. Disulfides may be cleaved 30 with dithiothreitol, glutathione, β -mercaptoethylamine or other conventional reagent. Cleavage of the disulfide may occur prior to or after conjugation to the polypeptide.

Depending upon the particular metal, various conditions and techniques will be employed for preparing the metal chelate. To prepare the technetium chelate, the chelating compound as carboxylate or activated ester is combined with a pertechnetate solution in the presence of a reducing agent, e.g., stannous or 35 dithionite under conventional conditions, whereby the technetium chelate is formed as a stable salt. The rhenium chelate may be formed by reducing perrhenate with stannous ion in the presence of citrate and the N_2S_2 ligand. Yields are 50% or greater after 1 hr at 50 °C.

The chelated acid may be already esterified or esterified in accordance with conventional ways. If already esterified, a labile complex such as Tc-99m gluconate will be prepared which will allow exchange to 40 the N_2S_2 activated ester ligand forming a complex suitable for protein conjugation. Alternatively, the carboxylic acid may be activated by employing a water soluble carbodiimide, e.g., EDCI, in an aqueous medium in the presence of at least a stoichiometric amount, preferably an excess of the hydroxylic compound. A suitably buffered aqueous medium may be employed. Excess carbodiimide can be converted to urea by adding acetate. The aqueous medium may then be used directly without further purification for 45 conjugation to the polypeptide. Desirably, the polypeptide will be added to the ester containing aqueous medium at a convenient concentration at a mildly alkaline pH, generally in excess of about 7.5 and less than about 9 and the reaction allowed to proceed for a sufficient time for all of the active ester to either react with the polypeptide or be substantially completely hydrolyzed. Usually, the time will be less than about 6 hr and more than about 30 min, with temperatures ranging from about 0 to 50 °C, usually not 50 exceeding about 40 °C. The particular conditions will be selected in accordance with the particular activated ester, the pH, the activity of the polypeptide, and the like.

It is also feasible but less preferable to conjugate the chelating agent (N_2S_2) to the polypeptide in the absence of the metal ion. The carboxylic acid group would be linked to the polypeptide to form a stable covalent link, e.g., an amide linkage, followed by the addition of the metal in a reduced, chelated, 55 exchangeable form. As chelates, α - or β -dioxo compounds are useful. Conveniently, the metal ion could be added as a weakly chelated ion or in the presence of a weakly chelating group, such as a uronate, e.g. gluconate.

The subject chelates will be administered to the mammalian host, normally by injection, intravenously,

EP 0 188 256 B1

intra-arterially, peritoneally, intratumorally, or the like, depending upon the particular site at which the radionuclide is desired. Generally, from about 0.1 to 2ml will be injected into a host, depending upon the size of the host, with about 0.001 to 50 μ Ci/kg of host. For human hosts the dosage will usually be about 10-50mCi/70kg host, more usually about 25-35mCi/70kg host. For lower mammals, e.g. mice, 1 μ Ci for
 5 biodistribution studies, while up to or greater than 500 μ Ci for imaging studies. After administration of the radionuclide, depending upon its purpose, the host may be treated in various ways for detection or therapy.

The following examples are offered by way of illustration and not by way of limitation.

EXPERIMENTAL

10

Example 1

Synthesis of N,N'-bis(benzoylmercaptoacetyl)-3,4-diamino butyrate.

15 In a dry flask under nitrogen is placed 1.54g (0.010mole) of 3,4-diaminobutyric acid hydrochloride and 250ml of absolute ethanol. Dry HCl gas is then bubbled into the solution. The mixture is refluxed for one to two days until formation of the ethyl ester is complete. The product is then concentrated to a dry solid and the hydrochloride ester dissolved by rapid stirring at ice bath temperature in a mixture of 50ml toluene and 50ml of saturated sodium bicarbonate. To this solution is added 5.0g (0.044mole) of chloroacetyl chloride in
 20 10ml of toluene by dropwise addition. After addition is complete, the mixture is allowed to come to room temperature and stirred for an additional 30min. Layers are separated and the aqueous portion is extracted twice with ethyl acetate. The organic layers are combined, washed with water and brine, and dried (magnesium sulfate). Removal of the solvent leaves the product as a white solid, which may be used without further purification.

25 A solution of 1.41g (about 4.45mmole) of the bis-chloroacetamide is prepared in 10ml of dry ethanol under nitrogen. To this is added a solution of sodium thiobenzoate in dry ethanol (prepared from sodium methoxide (0.204g of sodium, 8.87mmole, and ethanol) which is reacted with 1.23g (8.90mmole) of thiobenzoic acid). After a few minutes at room temperature, precipitation occurs. The reaction is heated to reflux for 30min. It is then allowed to cool, diluted with ethyl acetate, washed with water and brine and dried
 30 (magnesium sulfate). Removal of solvent leaves a cream-colored solid which may be recrystallized from toluene.

Example 2

35 Radiolabeling with Tc-99m.

1. The product prepared in Example 1 (0.1mg) is dissolved in 0.3ml of ethanol by heating and 30 μ l of 5N sodium hydroxide and 0.3ml of water added in succession. After heating for 15min at 95 $^{\circ}$ C during which time the ethanol evaporated, an essentially aqueous solution of the hydrolyzed ligand is left. To the
 40 mixture is then added generator pertechnetate in saline (0.5ml or less) which includes about 30mCi or less of Tc-99m and 0.5mg of freshly dissolved sodium dithionite; or (2) after allowing the mixture to stand for a short period at room temperature, the mixture is heated to 95 $^{\circ}$ C for an additional 15min and the pH adjusted to about 8.

2. The protected thiol, free carboxylic acid ligand of Example 1, 0.10mg, is added to 20mg of sodium gluconate and 0.010mg of SnCl₂·2H₂O, pH adjusted to 5. The Tc-99m as pertechnetate is added to the
 45 mixture and the mixture heated at 95 $^{\circ}$ C for 5 min.

The product mixture may be purified by preparative HPLC, using a 25cm octadecylsilane column (Altex Model 312 chromatograph; 4.6x250mm ODS ultrasphere, 5 μ) and eluting with 95% 0.01M sodium phosphate (pH 6) and 5% ethanol with a flow rate of 1.0ml/min. The preparations are analyzed for reduced
 50 hydrolyzed technetium on silica gel thin-layer strips.

Example 3

Formation of activated esters.

55

The conditions for formation of the activated esters are as follows: Into a reaction flask is introduced the carboxylic acid ligand or tracer level of metal complex carboxylate and an equimolar amount of the hydroxylic compound and a small excess, about 25% excess, of 1-ethyl-3-dimethylaminopropyl car-

EP 0 188 256 B1

bodilimide hydrochloride (ECDI) and 400 μ l of dimethylformamide (DMF). Upon completion of the reaction, sodium acetate is added to quench unreacted ECDI and the solution is ready for use for conjugation.

The protein to be conjugated is dissolved in 0.2M borate buffer, pH 8.5 to 9.0, to a protein concentration of about 2 to 5mg/ml. The mixture is allowed to stand at 4 °C until all of the protein has dissolved. To the aqueous protein solution at a pH adjusted to 8.5-9 is added the ester solution and the pH readjusted if necessary. The resulting conjugate is then preparatively chromatographed on an HPLC gel filtration column with 0.05M phosphate, pH 7.4, buffer as eluant.

In the following study, various conditions were employed, employing activated esters of technetium chelate prepared as described above for reaction with immunoglobulin under varying conditions of time, temperature, concentration, and pH. The following Table 1 indicates the results.

TABLE 1: Reactions of Activated Esters of N,N'-bis(mercaptoacetyl)-3,4-diaminobutanoate (Tc-99m) with Immunoglobulin under Different Conditions

Activated Ester	pH	Protein mg/ml	T(°C)	t(min)	% Labeled Protein	% Ester Hydrolyzed	% Ester Unreacted
p-nitrophenyl	6.94	1.0	23	22	0.5	3.7	95.8
	6.94	1.0	23	77	1.7	8.2	90.1
	6.94	1.0	37	60	4.3	17.5	78.2
	6.94	1.0	37	105	6.4	24.9	68.7
	8.70	1.0	23	105	12.0	33.0	68.7
	8.70	1.0	34	120	15.0	69.0	8.0
	8.70	1.0	23	320	20.0	50.0	23.0
	9.69	1.0	23	120	13.0	76.0	7.0
2-chloro-4-nitrophenyl	8.55	1.0	23	240	34.0	62.0	4.0
	8.55	1.0	0	240	37.0	60.0	3.0
	9.22	1.0	23	300	26.0	72.0	2.0
	9.22	1.0	0	330	30.0	66.0	4.0
	8.61	3.0	23	30	46.0	39.0	15.0
	8.61	3.0	0	60	38.0	46.0	16.0
	8.61	3.0	23	75	51.0	38.0	11.0
	8.61	3.0	0	100	44.0	42.0	14.0
hydroxybenzotriazole	6.80	1.0	23	30	8.4	45.4	46.0
	6.80	1.0	23	60	10.3	48.3	39.0
	6.80	1.0	23	120	10.8	53.3	35.8
	8.50	1.0	23	90	10.3	61.0	23.0
	8.70	1.0	5	120	32.0	30.0	32.0
	8.70	1.0	5	240	34.0	30.0	30.0
	9.25	1.0	5	45	12.0	33.0	53.0
	9.25	1.0	5	90	13.0	32.0	53.0
	9.40	1.0	23	60	4.3	66.0	21.0

EP 0 188 256 B1

Example 4

Synthesis of 4,5-diaminopentanoate.

5 To a solution of 50.5g of sodium bicarbonate in 200ml of water was added 85.0g of glutamic acid gamma-ethyl ester and the mixture cooled in an ice-salt bath. While maintaining the temperature between 0-5°C, 40g of carbobenzoxy chloride was added and the mixture stirred for 5hr followed by warming to room temperature and stirring for an additional 2hr. After extraction 2x100ml of ether, the mixture was acidified with 6N HCl to Congo red (pH 3). The separated oil was extracted with 3x100ml methylene
10 dichloride, the combined organic layers washed with brine and water and then dried over anh. sodium sulfate. Evaporation and crystallization from 200ml carbon tetrachloride gave a yield of 46.3g (77%) MP86-88°C.

To a solution of 46g of the above product in 45ml of THF at 35-40°C was rapidly added BH₃-THF (0.18mmol in 178ml). After 3hr, an aliquot on TLC (ethyl acetate hexane 4:1) showed substantially complete
15 conversion to the alcohol.

Fifty ml of ethanol was added to the reaction mixture and the mixture evaporated to dryness. After repeating the procedure twice with 100ml of ethanol, the residue was suspended in water, extracted with ethyl acetate and the organic layer washed successively with 2x100ml of 2% aqueous bicarbonate and water, followed by drying over anh. sodium sulfate. The organic solvent was then evaporated, the residue
20 dissolved in hexane and upon cooling gave 30.8g (71%) yield of a low-melting solid. MP86-88°C; TLC (R_f ethyl acetate-hexane 0.19).

The alcohol (29.5g) prepared above was dissolved in 90ml of pyridine (0-5°C) and 19.5g of tosyl chloride added at once. Precipitation of pyridiniumhydrochloride was observed after 1hr and the mixture stirred for 2hr more, followed by storage at 4°C overnight. The solution was poured with stirring into a liter of ice-water and the resulting solid isolated by filtration, washed with water and dried in a desiccator
25 overnight to yield 35g (80%) of the tosyl ester. MP73-76°C.

To the tosyl ester (22.45g) in 150ml of DMF was added 3.9g of sodium azide and the mixture heated at 50-55°C for 3hr. At the end of this time, the DMF was removed in vacuo at 5-10 torr., cold water added and filtered. The resulting azide was dried in a desiccator overnight to yield 14.56 (91%) of the desired
30 product. MP60-63°C.

Into 227ml of 1N HCl-ethanol (abs) was dissolved 14g of the above azide and the solution carefully added to 1.4g of platinum oxide in a hydrogenation bottle. The mixture was hydrogenated at 50-55°C for 48hr and the course of the reduction followed by TLC. At completion of the reaction, the catalyst was removed by filtration, the filtrate evaporated to dryness and the residue dissolved in 325ml of 6N HCl and
35 the mixture refluxed for 36hr. After filtration and evaporation to dryness, the residue was dissolved in 100ml of water, the water evaporated and the process repeated twice. The residue was triturated with ethanol to yield 8.3g (91%) of the diamino acid product. MP>250°C.

Example 5

40 Synthesis of Antibody N₂S₂ Conjugate Using o-Nitrophenyl Disulfide Protected Ligand.

To 2.05g of the above diamino acid dissolved in 50ml of DMF was added triethylamine (3ml) and succinimidyl S-benzoyl thioglycolate (5.86g) and the mixture stirred for 15min. Dimethylformamide was
45 removed under vacuum and 100mL of cold water was added. The precipitated oil solidified on standing. The solid was filtered, dried and crystallized from ethyl acetate. MP126-127°C.

To sodium ethoxide (140mg sodium) in 30ml ethanol was added 0.966 of the above product and the mixture stirred overnight at room temperature. After evaporating the solvent in vacuo, the residue was dissolved in glacial acetic acid, the solvent evaporated and the process repeated twice. The residue was
50 redissolved in 30ml of glacial acetic acid and 0.77g of o-nitrophenylsulfenyl chloride added and the mixture stirred at room temperature for 24hr. The reaction was monitored by TLC (acetonitrile-water 95:5) and at completion of the reaction, the acetic acid was removed in vacuo and cold water added. The solid precipitate was filtered, washed with cold ethanol (10-15ml) and dried in vacuo for 12hr over P₂O₅. The yield was 1.03g (88%). MP>200°C. TLC:acetonitrile:water 95:5 R_f 0.39.

55 To the bis-(di-o-nitrophenyl)disulfide (0.293g) suspended in 50ml THF (anh.) was added N-hydroxysuccinimide (63mg) followed by dicyclohexylcarbodiimide (113mg) and the mixture stirred for 48hr at room temperature. The solution was concentrated to about 15-20ml and cooled, the precipitate removed by filtration and the filtrate diluted with 25-30ml of ethyl acetate, followed by washing the organic layer with

EP 0 188 256 B1

water. The organic layer was dried over magnesium sulfate, concentrated to 20ml and cooled. The resulting precipitate was filtered, the filtrate concentrated to about 10ml and cooled to about 10°-15° C. After filtration, the filtrate was maintained at about 4° C for 2-3hr. Addition of anh. ether to the cold solution resulted in a yellow precipitate (about 95mg), followed by a second crop of about 90mg of an impure product.

The antibody conjugation reaction was contained in a final volume of 40ml: 1.8mg (1.72×10^{-5} moles) bis-(di-o-nitrophenyl)disulfide) N_2S_2 ligand, 178mg of mouse monoclonal antibody (IgG, 1.2×10^{-6} moles), 4.0ml of redistilled DMF, 0.05M sodium borate buffer pH 8.5. After stirring 90min at room temperature, 4.4ml of 5N sodium chloride and 1.9ml of 100mM dithiothreitol were added. After an additional 30min the reaction mixture was centrifuged to remove any particulates and the supernatant fractionated by gel filtration column chromatography. The column eluent was monitored at 280nm and the fractions containing the monomeric antibody conjugate were pooled and concentrated in an Amicon stirred cell (30,000 molecular weight cutoff). Final yield was 141mg (82%).

Example 6

Technetium-99m Labeling of Antibody-Ligand Conjugate with Tc-Tartrate.

Stannous tartrate kits were prepared from degassed solutions of 0.5ml disodium tartrate (150mg/ml) and 0.1ml stannous chloride, (1.0mg/ml in ethanol) in an evacuated vial under nitrogen atmosphere. To a stannous tartrate kit, sodium pertechnetate 0.5ml (~15mCi) was added and heated at 50° C for 10-15min. After cooling to room temperature, quality control for Tc-99m tartrate and insoluble Tc-99m was carried out on Gelman ITLC using methyl ethyl ketone and 0.01M sodium tartrate pH 7.0 eluents, respectively. Tc-99m tartrate formation was typically 98-99% with insoluble Tc-99m values ranging from 0.1 to 0.2%.

In an evacuated vial, 100μl saline, 200μl of sodium phosphate (0.2M, pH 8.0) and 200μl of antibody-ligand conjugate (1.9mg/ml) were added successively. Immediately after adding the conjugate, 250μl of Tc-99m tartrate (~3 to 5mCi was added and heated at 50° C for 1hr. Percent technetium bound to protein and the formation of pertechnetate were determined by ITLC using 50% MeOH:10% ammonium acetate (1:1) and 1-butanol eluents, respectively. Technetium incorporation typically ranged from 70-88% on a ligand-Ab conjugate with a ligand per antibody ratio of 1.5 to 3.0.

TABLE 2: Comparative Biodistribution of Tc-99m and Iodine-125 Anti-melanoma Antibody 9.2.27 in Mice Bearing Melanoma Tumors From FEMX Cell Line.

Organ	Tumor	Liver	Spleen	Lung	Stomach	Thyroid	Kidney
Tc-99m	5.78* ±0.32	1.54 ±0.19	1.34 ±0.14	1.79 ±0.67	0.26 ±0.15	0.61 ±0.05	1.72 ±0.12
I-125	3.97 ±0.61	1.07 ±0.17	1.59 ±0.03	1.81 ±0.12	2.99 ±1.89	7.79 ±4.50	1.33 ±0.04

* Data are mean ±S.D. percent injected dose per gram for three mice at 48hr post injection.

The method of Hwang et al., Cancer Res. (1985) 45:4150-4155 was employed.

Example 7

Labeling of Antibody with Preformed Tc-99m Pentanoyl N_2S_2 Chelate.

EP 0 188 256 B1

A Tc-99m chelated derivative was conjugated to an antibody as follows. Tc-99m (75mCi) chelated by N,N'-bismercaptoacetyl 4,5-diaminopentanoic acid was prepared by dithionite reduction of Tc-99m pertechnetate at basic pH with 25 μ g of the N₂S₂ ligand. The acid was activated by adding the above complex at pH 7 in 0.5ml water to 100 μ l of water:acetonitrile (1:9) containing 3.0mg of 2,3,5,6-tetrafluorophenol and 100 μ l of H₂O:acetonitrile (1:9) containing 7.5mg of 1-cyclohexyl-3-(2-morpholinoethyl)carbodiimide(morpho CDI) added. After storing for 18hr at room temperature, the mixture was purified using a Baker-10 SPE reversed phase C₁₈ column. The column was conditioned with 2ml of ethanol followed by washing with HPLC grade water. The reaction mixture was then added to the column, the column washed 4x with 2ml volumes of 10% methanol in 0.01M sodium phosphate, pH 7.0 and the ester complex eluted with 2.5ml portions of acetonitrile. The first eluent contained 8.5mCi and the second 0.18mCi. The yield was 86% after accounting for decay.

To a 2ml vial was added 4.5 mCi of activated ester complex in acetonitrile, the solvent evaporated in a nitrogen stream, and 0.40ml of sodium borate (0.5M, pH 9.0) added. With stirring, 30 μ l (9.14mg/ml) of anti-melanoma antibody (9.2.27) was added. The final protein concentration was 0.52mg/ml. The reaction was followed with TLC using Gelman ITLC SG strips and eluting with 50% aqueous methanol:10% ammonium acetate (1:1), indicating that 47% protein bound Tc-99m at 15min and 59% at 30min at room temperature. The Tc-99m labeled protein was purified by Centricon-10k filter centrifugation. A sample of 92.4% protein bound Tc-99m showed 84.0% binding to FEMX melanoma cells.

20 Example 8

Preparation of Re-186 4,5-dimercaptoacetamidopentanoyl-antibody (anti-melanoma antibody 9.2.27).

In an evacuated vial is combined 100 μ l of H₂O, 100 μ l acetonitrile, 100 μ l of citrate solution (28.8mg, 1.5x10⁻⁴ mol), 50 μ l of ligand (tetrafluorophenyl 4,5-di-(tetrahydropyranylmecaptoacetamido)pentanoate (0.40mg; 6.5x10⁻⁷ mol), 50 μ l of stannous chloride (0.5mg, 2.6x10⁻⁶ mol), and 50 μ l of Re-186 perrhenate in acetonitrile (4.25 μ g, 2.3x10⁻⁸ mol). The mixture is heated at 50 °C for 1hr and then 0.30ml of 1N NaOH is added.

The tetrafluorophenyl ester product of the Re-186 N₂S₂ complex is purified on a C₁₈ Baker-10 SPE column. After application to the column, impurities are washed off with 2x3ml of H₂O and 4x3ml of 10% CH₃OH/0.01M phosphate, pH 7. The product is eluted with 2ml of acetonitrile and then the solution is reduced to dryness under a stream of nitrogen. Yields of product are about 60%.

Conjugation of Re-186 N₂S₂ complex is done by addition of antibody (160 μ l of 5mg/ml) (Morgan et al., Hybridoma (1981) 1:27), in 340 μ l of borate buffer (0.5M, pH 9). After 30min at room temperature, 58% of the radioactivity was protein bound. Immunoreactivity determined by binding of radioactivity to FEMX melanoma cells was 80% after correction for nonprotein bound material.

Example 9

40 Synthesis of Imidate form of N₂S₂ Ligand,
Conjugation to Antibody and Radiolabeling with Tc-99m.

2,3-(Bis-carbobenzyloxy)diaminopropan-1-ol [2]

45 A 500mL hydrogenation bottle was charged with 55g (0.25 mol) of 2,3-dibromopropanol (Aldrich) and 300mL of 28-30% aqueous NH₄OH solution. The mixture was stoppered with an internal thermometer and heated to 75-85 °C while shaking on a Parr shaker for 23hr. When cool, shaking was stopped and the mixture was carefully opened. The mixture was evaporated to a volume of 50mL by passing N₂ gas through it while heating on an oil bath. While hot, 50mL of EtOH was added and the mixture was allowed to cool. 50 The hydrogen bromide salt of 2,3-diaminopropan-1-ol was collected by filtration and dried in vacuo to yield 50g of hard chunks of white solid which was used without further purification.

A solution of 25g of the crude salt in 110mL of 4N NaOH was cooled to 0 ° (ice bath), and to the solution was added a solution of 31.4mL (0.22mol, 37.5g) of benzylchloroformate in 100mL of CH₂Cl₂. The mixture was stirred rapidly for 30min at 0 ° and 16hr at room temperature. The CH₂Cl₂ phase was collected, washed with 75mL of brine, dried (MgSO₄), filtered and concentrated. The resulting solid was washed with 100mL of Et₂O, collected by filtration, and dried under vacuum to give 10.7g (24%) of 2 as a white solid which could be recrystallized from CHCl₃/hexane to give tiny needles. MP119-120 °C.

EP 0 188 256 B1

2,3-(Bis-carbobenzyloxy)(diaminopropyl-1-methanesulfonate) [3]

To a suspension of 10.68g (30mmol) of 2 and 6.27mL (4.55g, 45mmol) of Et₃N in 150mL of CH₂Cl₂ cooled to 0° under N₂ atmosphere was added 2.55mL (3.78g, 33mmol) of methanesulfonyl chloride, and the mixture was stirred for 30min at 0° C. The resulting clear solution was washed successively with 75mL of 5% HCl, 75mL of H₂O, 75mL of 5% NaHCO₃, and 75mL of sat. aq. NaCl (all chilled in ice). The CH₂Cl₂ phase was dried (MgSO₄), filtered, concentrated, and crystallized from CHCl₃/hexane to yield 12.33g (94%) of white crystals. MP92-93° C.

3,4-(Bis-carbobenzyloxy)diaminobutyronitrile [4]

A mixture of 6.56g (15mmol) of 3, 1.08g (16.5mmol) of KCN, 0.40g (1.5mmol) of 18-crown-6, and 75mL of anhydrous acetonitrile (stored over 3A molecular sieves) was refluxed in a nitrogen atmosphere for 19hr. When cool, the mixture was partitioned between 100mL of 10% NaHCO₃ solution and 200mL of CH₂Cl₂. The CH₂Cl₂ layer was washed successively with 100mL portions of 5% HCl, water and brine. The CH₂Cl₂ phase was dried (MgSO₄), filtered, and concentrated to give 5.47g of brown oil. Two recrystallizations from CHCl₃/hexane yielded 2.68g of 4 as a white solid. MP111-112° C.

3,4-Diaminobutyronitrile dihydrogeniodide salt [5]

To 3.38g (13.3mmol) of I₂ in a 100mL flask under N₂ atmosphere was added 5.42mL (3.87g, 26.5mmol) of hexamethyldisilane. The mixture was immersed in a 45-50° C oil bath until solid I₂ dissolved (30min). The temperature was raised to 100° and held for 5min until the color disappeared. The solution was cooled to 0° with an ice bath and diluted with 13.3mL of CH₂Cl₂. To the 0° C solution was added dropwise over 5min a solution of 1.96g (5.3mmol) of 4 in 13.3mL of CH₂Cl₂. The cooling bath was removed and the mixture was stirred in the dark for 3hr at room temperature. To the mixture was added 2.15mL (1.70g, 53mmol) of MeOH and stirring was continued overnight (16hr). The mixture was cooled to 0° C and the solid was collected by filtration and dried under vacuum to give 1.75g (100%) of a tan solid 5 which was characterized as its dibenzoyl derivative.

3,4-Dibenzoylmercaptoacetamidobutyronitrile [6]

To a mixture of 3.27g (10mmol) of 5, 7.33g (25mmol) of N-succinimidyl S-benzoylmercaptoacetate and 10mL of DMF was added at 0° C under N₂ atmosphere, 3.48mL (2.52g, 25mmol) of triethyl amine. The cooling bath was removed and the mixture was stirred for 1hr. The mixture was diluted with 50mL of 5% HCl solution and extracted with 2x50mL of CH₂Cl₂. The combined CH₂Cl₂ phases were washed with 100mL of 5% NaHCO₃ solution, dried (MgSO₄), filtered, and concentrated in vacuo to yield 6.75g of a purple tinted solid.

Purification was accomplished by chromatography (silica gel, EtOAc) and crystallization of purified fractions (CHCl₃/hexane) to give 3.20g (70%) of white solid. Mp125-127° C.

3,4-Bis-methyldithioacetamidobutyronitrile [7]

To a suspension of 455mg (1.0mmol) of 6 in 6mL of EtOH at room temperature under N₂ atmosphere was added 2.2mL of 1N aqueous NaOH. The mixture was stirred at room temperature for 1.6hr, and to the resulting clear solution was added 226μl of methyl methanethiolsulfonate. The mixture was stirred for 3hr and partitioned between 20mL of pH 7 buffer solution and 2x20mL of CH₂Cl₂. The combined aqueous layers were dried (MgSO₄), filtered, and concentrated to give 591mg of pink residue. Purification by silica gel chromatography (EtOAc) and crystallization from CHCl₃/hexane gave a total of 217mg (64%) of white amorphous solid 7. MP121-123° C.

Methyl 3,4-bis-methyldithioacetamidobutyrimidate hydrogen chloride salt [1]

A suspension of 141mg (0.41mmol) of 7 in 1.66mL of MeOH and 4.15mL of Et₂O was cooled to -20° C (CO₂/CCl₄), and HCl gas was passed through the mixture via septum inlet for 5min, until most of the solids had dissolved and the solution was saturated with HCl. The mixture was placed in the freezer in a desiccator for 66hr, and then concentrated under vacuum to produce a white foamy solid. The solid was broken up, washed with three portions of anhydrous Et₂O, dried under vacuum to give 111mg (66%) of 1 as

EP 0 188 256 B1

a off-white solid which decomposed on heating and also decomposed after several days in a freezer.

Preparation of Antibody - Methyl 3,4-bis-methyldithioacetamidobutyrimidate Conjugate

5 A 2mg/ml stock solution of the N_2S_2 ligand was prepared in dry acetonitrile. The solution was standardized by determining the disulfide content using 2-nitro-5-thiosulfobenzoate (Thannhauser et al., Anal. Biochem. (1984) 138:181) and the ligand concentration was found to be 5.30mM.

For conjugation to mouse monoclonal antibody, 0.16ml of N_2S_2 ligand-acetonitrile solution was added to the reaction vial and the solvent removed with a stream of dry nitrogen. Antibody (0.62ml of 8.1mg/ml
10 solution) and 1.0ml of 0.2M sodium bicarbonate buffer pH 9.5 were mixed and then added to the reaction vessel containing the dried ligand. After stirring 30min at room temperature, the entire solution was added to a fresh vial containing the same amount of dried ligand and the solution stirred another 30min. The conjugated antibody was purified by Sephadex G-25 chromatography in 50mM sodium phosphate pH 7.5, 0.5M sodium chloride. The protein containing fractions were pooled and concentrated in an Amicon stirred
15 cell to a concentration of about 2mg/ml. The solution was made 50mM in glutathione, stirred 25min, then purified by Sephadex G-25 gel filtration and concentrated as before. The final solution (1.7mg/ml) was stored at 4° C until use.

Radiolabeling of Antibody - Methyl 3,4-bis-methyldithioacetamidobutyrimidate Conjugate

20 Tc-99m tartrate was prepared in a total volume of 1.1ml of degassed water with 100 μ g $SnCl_2$, 9% (v/v) ethanol, 75mg disodium tartrate, and 3.2mCi sodium (Tc-99m) pertechnetate. The solution was heated at 50° C for 15min. To a separate vial was added 100 μ l of the Tc-99m tartrate solution, 100 μ l of 0.2M sodium bicarbonate, pH 10, and 100 μ g of the antibody conjugate. The total volume was then adjusted to 0.5ml with
25 0.15M sodium chloride and the solution incubated at 50° C for 60min. Analysis by HPLC (TSK column, 0.2M sodium phosphate pH 7.4, 0.15M sodium chloride) showed 95% of the Tc-99m was associated with the antibody conjugate.

Example 10

30 Preparation of S-terephthaloylsubstituted N_2S_2 Ligand.

The mono-tert-butyl ester of terephthalic acid 1 was prepared by the method of Buckle and Smith, J. Chem. Soc. (1971) 54:2821.

35 Succinimidyl ester 2 was prepared by stirring 1 with 1.2 molar equivalents of N-hydroxysuccinimide and 1.3 molar equivalents of 1,3-dicyclohexylcarbodiimide in dry THF at room temperature for 14-16hr. Thin layer chromatographic analysis indicated the reaction had gone to completion. The dicyclohexylurea was then removed by filtration and the resulting liquid was concentrated in vacuo to yield 2 as a white solid. Final purification of 2 was accomplished by flash chromatography.

40 The thioester 3 was prepared by dissolving 1.0 molar equivalents of mercaptoacetic acid and 2.0 molar equivalents of 4-dimethylaminopyridine in dry THF. The succinimidyl ester 2 was added to the stirring solution. After stirring for 5hr the reaction was complete as indicated by thin layer chromatographic analysis. The THF was removed in vacuo and the residue was dissolved in CH_2Cl_2 . The solution was then washed with dilute aqueous HCl and dried over anhydrous $MgSO_4$. Filtration and evaporation of the solvent gave 3
45 as a colorless oil which solidified upon standing.

The succinimidyl ester 4 was prepared by the method of Subramanian (R.F. Schnelder et al., J. Nucl. Med., (1984) 25:223-229).

The carboxylic acid 5 was prepared by dissolving 4,5-diaminopentanoic acid dihydrochloride salt in 1:4 $H_2O:CH_3CN$ containing 3.0 molar equivalents of triethylamine and subsequently adding 2.0 molar equivalents of the succinimidyl ester 4. After stirring for 14-18hr at room temperature, TLC analysis showed the reaction to be complete and the solvent was removed in vacuo. The residue was dissolved in ethyl acetate and washed with dilute aqueous HCl, water, and brine. The ethyl acetate layer was then dried over anhydrous Na_2SO_4 . After filtration and removal of the solvent, a waxy solid was obtained which was recrystallized from a mixture of ethyl acetate and hexane to give 5 as a white solid.

55 Tetrafluorophenyl ester 6A was prepared by dissolving 5, along with 1.2 molar equivalents of 2,3,5,6-tetrafluorophenol in dry THF. 1,3-dicyclohexylcarbodiimide (1.2 molar equivalents) was added to the mixture and the mixture was stirred for 12-15hr. Analysis by thin layer chromatography indicated the reaction was complete. The dicyclohexylurea was removed and by filtration and the solvent was removed in vacuo. The

EP 0 188 256 B1

residue was purified by flash chromatography to yield the ester 6A as a white solid.

Succinimidyl ester 6B was prepared by dissolving 5 along with 1.2 molar equivalents of N-hydroxysuccinimide in dry THF. 1,3-dicyclohexylcarbodiimide (1.2 molar equivalents) was added to the mixture and the mixture was stirred at room temperature for 14-18hr. Thin layer chromatographic analysis indicated the reaction had gone to completion. The dicyclohexylurea was removed by filtration and the solvent was removed in vacuo. The residue was dissolved in ethyl acetate and washed with water. The ethyl acetate solution was dried over anhydrous Na_2SO_4 . The drying agent was removed by filtration and the solvent was removed in vacuo. The resulting residue was purified by flash chromatography to yield the succinimidyl ester 6B as a white solid.

Removal of the tert-butyl protecting groups was accomplished by dissolving the tetrafluorophenyl ester 6A in CH_2Cl_2 and treating the solution with excess trifluoroacetic acid, initially at 0°C , then stirring to room temperature for 3hr. Thin layer chromatographic analysis showed that the reaction had gone to completion. The solvent and excess trifluoroacetic acid were then removed in vacuo to yield a white to colorless solid which was recrystallized from $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ to give 7A as a white powder.

In the case of the succinimidyl ester 6B, the tert-butyl protecting groups were removed as described above for compound 6A. It was necessary, however, to purify the product 7B by flash chromatography.

These reaction sequences were also carried out starting with the mono tert-butyl ester of isophthalic acid to obtain the analogous meta isomers of the products described above.

Example 11

Conjugation of N-hydroxysuccinimidyl 4,5-diterephthaloylmercaptoacetamidopentanoate to IgG antibody.

The conjugation was carried out in a total volume of 2.0ml and contained $480\mu\text{g}$ (7.1×10^{-7} moles) N_2S_2 ligand active ester 6B, 0.2ml redistilled DMF (10%), 0.15M sodium chloride, 0.05M sodium borate, pH 8.5, and 10.0mg mouse monoclonal antibody (6.7×10^{-8} moles). After stirring 90min at room temperature the reaction was fractionated by gel filtration over Sephadex G-2B in 0.05M sodium phosphate buffer pH 7.5 with 0.15M sodium chloride. The excluded volume containing the conjugated antibody was collected. To remove any residual non-protein material the conjugate was dialyzed 18hr against 0.05M sodium phosphate, pH 7.5, with 0.15M sodium chloride. Final yield of protein was 100%.

Example 12

Tc-99m Labeling of 4,5-diterephthalylmercaptoacetamidopentanoyl-IgG Antibody Conjugate.

To $120\mu\text{l}$ saline $200\mu\text{l}$ of 0.2M sodium phosphate buffer, pH 8, and $80\mu\text{l}$ of the terephthaloyl sulfur protected N_2S_2 conjugate (4.66mg/ml), $250\mu\text{l}$ of the Tc-99m tartrate ($\sim 4\text{mCi}$) prepared as previously described was added. The reaction mixture was heated at 50°C for 1hr which resulted in a Tc-uptake of 90%.

Following the above procedure the isophthaloyl analog could also be prepared.


It is important that the resulting product provide for maximum formation of the radionuclide conjugates. In addition, there is the concern about the time, since the radioisotopes do decay with time. Thus, by using the compounds of the subject invention, one can rapidly conjugate proteins to provide radionuclide substituted reagents for use in vivo. The reagents can be provided in pure form, good yield, and the radionuclide metal is stably maintained as a chelate with the protein for use in vivo. Thus, one can safely direct the radionuclide to a desired site, where only low levels of radioactivity will be non-specifically directed and bound.

Claims

Claims for the following Contracting States: BE, CH, DE, FR, GB, IT, LI, LU, NL, SE:

1. A compound of the formula:

EP 0 188 256 B1



wherein:

15

M is a metal or metal oxide radionuclide ion capable of being chelated;

one of Z¹, Z², Z³, Z⁴, is R'CW'(HNv)_n, Y', and the others are H₂ or =O;

20

R' is an aliphatic divalent radical of from 1 to 6 carbon atoms and 0 to 2 heteroatoms;

Y' is the leaving group of an active ester, -NH₂, -NHNH₂, or a polypeptide of at least about 1000 molecular weight;

25

V' is of the formula R'CW';

W' is =NH or =O, with the proviso that the W' bonded to the carbon atom bonded to Y' is H₂ when Y' is -NH₂;

30

X' is a bond, methylene or CHZ⁴;

(A)'s are the same or different and are hydrogen or lower alkyl of from 1 to 6 carbon atoms; and

n' is 0 or 1.


35

2. A compound according to Claim 1, wherein Y' is a polypeptide.

3. A compound according to Claim 2, wherein Y' is an immunoglobulin or specific binding fragment thereof.

40

4. A compound of the formula:



wherein:

55

one of Z¹, Z², Z³, Z⁴, is RCW(HNv)_nY, and the others are H₂ or =O;

R is an divalent radical of from 1 to 6 carbon atoms and 0 to 2 heteroatoms;

EP 0 188 256 B1

W is =NH or =O, with the proviso that the W bonded to the carbon atom bonded to Y is H₂ when Y is -NH₂;

V is the same or different from RCW and comes within the definitions of RCW;

n is 0 or 1;

T is a removable protective group, hydrogen, or an alkali metal ion;

Y is an organic oxy compound forming an active ester, -NH₂, -NHNH₂, or a polypeptide of at least two amino acids;

X is a bond, methylene or CH₂ⁿ; and

the A's are the same or different and are hydrogen or lower alkyl of from 1 to 6 carbon atoms.

5. A compound according to Claim 4, wherein X is a bond.

6. A compound according to Claims 4 or 5, wherein Y is an activated phenoxy group.

7. A method for preparing a radiolabeled polypeptide which comprises:

combining in an aqueous medium a polypeptide and a compound of the formula according to Claim 1, wherein Y is the leaving group of an active ester capable of forming an amide or amidine link with said polypeptide, or NH₂ or NHNH₂, with the proviso that when Y is NHNH₂ or -NH₂, said polypeptide has an oxo-group as a result of glycol cleavage of a sugar bound to said polypeptide and said combining is under reductive amination conditions;

wherein said compound forms a stable covalent link to said polypeptide with the substantial absence of unchelated radionuclide.


8. A method for preparing a radiolabeled polypeptide which comprises:

combining in an aqueous medium a compound according to anyone of Claims 4 to 6, wherein T is an alkali metal ion or a sulfur protective group and Y is said polypeptide, and a metal radionuclide in a reduced chelated exchangeable form;

whereby said radionuclide becomes chelated by said compound; and

washing said radionuclide containing compound to remove unchelated radionuclide bound to said compound.


9. A compound of the formula



in which T is a sulfur protective group.

EP 0 188 256 B1


10. A compound of the formula:



wherein M represents a metal radionuclide selected from Tc-99m and Re-186.

Claims for the following Contracting State: AT:

1. A method of making a compound of the formula:



wherein:

M is a metal or metal oxide radionuclide ion capable of being chelated;

one of Z¹, Z², Z³, Z⁴, is R'CW'(HNV)_n, Y', and the others are H₂ or =O;

R' is an aliphatic divalent radical of from 1 to 6 carbon atoms and 0 to 2 heteroatoms;

Y' is the leaving group of an active ester, -NH₂, -NHNH₂, or a polypeptide of at least about 1000 molecular weight;

V' is of the formula R'CW';

W' is =NH or =O, with the proviso that the W' bonded to the carbon atom bonded to Y' is H₂ when Y' is -NH₂;

X' is a bond, methylene or CHZ⁴;

(A')s are the same or different and are hydrogen or lower alkyl of from 1 to 6 carbon atoms; and

n' is 0 or 1.

comprising the step of chelating a metal or metal oxide radionuclide ion with a precursor of said compound.

2. A method according to Claim 1, wherein Y' is a polypeptide.

3. A method according to Claim 2, characterized in that Y' is an immunoglobulin or specific binding fragment thereof.

EP 0 188 256 B1


4. A method for preparing a radiolabeled polypeptide which comprises:

combining in an aqueous medium a polypeptide and a compound of the formula according to Claim 1, wherein Y is the leaving group of an active ester capable of forming an amide or amidine link with said polypeptide, or NH_2 or NHNH_2 , with the proviso that when Y is NHNH_2 or $-\text{NH}_2$, said polypeptide has an oxo-group as a result of glycol cleavage of a sugar bound to said polypeptide and said combining is under reductive amination conditions;

wherein said compound forms a stable covalent link to said polypeptide with the substantial absence of unchelated radionuclide.

5. A method for preparing a radiolabeled polypeptide which comprises: washing a radionuclide containing compound obtained according to Claim 1 to remove unchelated radionuclide bound to said compound.

6. A method according to anyone of the preceding claims, characterized in that a compound of the formula:




is prepared,

wherein M represents a metal radionuclide selected from Tc-99m and Re-186.

Revendications

REVENDEICATIONS pour les Etats contractants suivants: BE, CH, DE, FR, GB, IT, LI, LU, NL, SE.

1. Composé de formule



dans laquelle

M est un ion de radionucléide de métal ou d'oxyde métallique pouvant être chélaté;

l'un de Z^1 , Z^2 , Z^3 , Z^4 est $\text{R}'\text{CW}'(\text{HNV})_n\text{Y}'$ et les autres sont H_2 ou $=\text{O}$;

R' est un radical divalent aliphatique de 1 à 6 atomes de carbone et de 0 à 2 hétéroatomes;

Y' est le groupement labile d'un ester activé, $-\text{NH}_2$, $-\text{NHNH}_2$ ou un polypeptide de poids moléculaire d'au moins environ 1000;

V' a la formule $\text{R}'\text{CW}'$;

W' est $=\text{NH}$ ou $=\text{O}$, pourvu que le groupement W' lié à l'atome de carbone lié à Y' soit H_2 quand Y' est $-\text{NH}_2$;

X' est une liaison, un groupement méthylène ou CH_2 ;

les radicaux A' sont identiques ou différents et sont des atomes d'hydrogène ou des groupements alkyle inférieur ayant de 1 à 6 atomes de carbone; et


EP 0 188 256 B1

n' est 0 ou 1.

2. Composé selon la revendication 1, dans lequel Y' est un polypeptide.

3. Composé selon la revendication 2, dans lequel Y' est une immunoglobuline ou un fragment à liaison spécifique de celle-ci.

4. Composé de formule



dans laquelle

l'un de Z¹, Z², Z³, Z⁴ est RCW(HNV)_nY et les autres sont H₂ ou =O;

R est un radical divalent de 1 à 6 atomes de carbone et de 0 à 2 hétéroatomes;

W est =NH ou =O pourvu que le W lié à l'atome de carbone lié à Y soit H₂ quand Y est -NH₂;

V est identique à ou différent de RCW et prend les définitions de RCW;

n est 0 ou 1;

T est un groupement protecteur éliminable, un atome d'hydrogène ou un ion de métal alcalin;

Y est un groupement oxy organique formant un ester actif, -NH₂, -NHNH₂ ou un polypeptide d'au moins deux acides aminés;

X est une liaison ou un groupement méthylène ou CHZ⁴; et les radicaux A sont identiques ou différents et sont des atomes d'hydrogène ou des groupements alkyle inférieur de 1 à 6 atomes de carbone.

5. Composé selon la revendication 4, dans lequel X est une liaison.

6. Composé selon les revendications 4 ou 5, dans lequel Y est un groupement phénoxy activé.

7. Procédé de préparation d'un polypeptide à marqueur radioactif, qui consiste à combiner dans un milieu aqueux un polypeptide et un composé de formule selon la revendication 1, dans laquelle Y est le groupement labile d'un ester actif pouvant former une liaison amide ou amidine avec ledit polypeptide, ou -NH₂ ou -NHNH₂, pourvu que quand Y est -NHNH₂ ou -NH₂, ledit polypeptide possède un groupement oxo par suite de la coupure glycolique d'un sucre lié audit polypeptide et ladite combinaison se fasse dans des conditions d'amination réductrice; ledit composé formant une liaison covalente stable avec ledit polypeptide en l'absence substantielle de radionucléide non chélaté.

8. Procédé de préparation d'un polypeptide à marqueur radioactif, qui consiste :


à combiner dans un milieu aqueux un composé selon l'une quelconque des revendications 4 à 6, dans lequel T est un ion de métal alcalin ou un groupement protecteur du soufre et Y est ledit polypeptide, et un radionucléide métallique sous une forme échangeable chélatée réduite;

ledit radionucléide étant alors chélaté par ledit composé; et

à laver ledit composé contenant le radionucléide pour enlever le radionucléide non chélaté lié audit composé.


9. Composé de formule

EP 0 188 256 B1



15 dans laquelle T est un groupement protecteur du soufre.


10. Composé de formule



dans laquelle M représente un radionucléide de métal choisi entre Tc-99m et Re-186.

30 **REVENDICATIONS pour l'Etat contractant suivant: AT**

1. Procédé de préparation d'un composé de formule



45 M est un ion de radionucléide de métal ou d'oxyde métallique pouvant être chélaté;

l'un de Z^{1'}, Z^{2'}, Z^{3'}, Z^{4'} est R'CW'(HNV)_n, Y' et les autres sont H₂ ou =O;

R' est un radical divalent aliphatique de 1 à 6 atomes de carbone et de 0 à 2 hétéroatomes;

Y' est le groupement labile d'un ester activé, -NH₂, -NHNH₂ ou un polypeptide de poids moléculaire d'au moins environ 1000;

50 V' a la formule R'CW';

W' est =NH ou =O, pourvu que le groupement W' lié à l'atome de carbone lié à Y' soit H₂ quand Y' est -NH₂;

X' est une liaison, un groupement méthylène ou CHZ^{4'};


les radicaux A' sont identiques ou différents et sont des atomes d'hydrogène ou des groupements alkyle inférieur ayant de 1 à 6 atomes de carbone; et

55 n' est 0 ou 1

comprenant l'étape consistant à chélatier un ion de radionucléide de métal ou d'oxyde métallique avec un précurseur dudit composé.

EP 0 188 256 B1

2. Procédé selon la revendication 1, dans lequel Y' est un polypeptide.
3. Procédé selon la revendication 2, dans lequel Y' est une immunoglobuline ou un fragment à liaison spécifique de celle-ci.
- 5 4. Procédé de préparation d'un polypeptide à marqueur radioactif, qui consiste à combiner dans un milieu aqueux un polypeptide et un composé de formule selon la revendication 1, dans laquelle Y est le groupement labile d'un ester actif pouvant former une liaison amide ou amidine avec ledit polypeptide, ou -NH₂ ou -NHNH₂, pourvu que quand Y est -NHNH₂ ou -NH₂, ledit polypeptide possède un groupement oxo par suite de la coupure glycolique d'un sucre lié audit polypeptide et ladite combinaison se fasse dans des conditions d'amination réductrice; ledit composé formant une liaison covalente stable avec ledit polypeptide en l'absence substantielle de radionucléide non chélaté.
- 10 5. Procédé de préparation d'un polypeptide à marqueur radioactif, qui consiste : à laver un composé contenant un radionucléide obtenu selon la revendication 1 pour enlever le radionucléide non chélaté lié audit composé.
- 15 6. Procédé selon l'une quelconque des revendications précédentes, caractérisé en ce qu'on prépare un composé de formule
- 20




dans laquelle M représente un radionucléide de métal choisi entre Tc-99m et Re-186.

Patentansprüche

Patentansprüche für folgende Vertragsstaaten: BE, CH, DE, FR, GB, IT, LI, LU, NL, SE.

- 35 1. Verbindung mit der Formel:



50 in der

M ein chelatisierbares Metall- oder Metalloxid-Radionuklidion ist;

eine der Gruppen Z¹, Z², Z³ und Z⁴, R'CW(HNV)_n, Y' ist und die anderen H₂ oder = O sind;

55 R' ein aliphatischer, zweibindiger Rest mit 1 bis 6 Kohlenstoffatomen und 0 bis 2 Heteroatomen ist;

Y' die austretende Gruppe eines aktiven Esters, -NH₂, -NHNH₂ oder ein Polypeptid mit einem

EP 0 188 256 B1

Molekulargewicht von mindestens ungefähr 1000 ist;

V' die Formel R'CW' aufweist;

5 W' = NH oder =O ist, mit der Bedingung, daß das an das mit Y' verknüpfte Kohlenstoffatom gebundene W' H₂ ist, wenn Y' -NH₂ ist;

X' eine Bindung, Methylen oder CHZ^{4'} ist;


10 die Gruppen A', die gleich oder verschieden sein können, Wasserstoff oder niederes Alkyl mit 1 bis 6 Kohlenstoffatomen sind; und

n' 0 oder 1 ist.

15 2. Verbindung nach Anspruch 1, in der Y' ein Polypeptid ist.

3. Verbindung nach Anspruch 2, in der Y' ein Immunoglobulin oder spezifisches bindendes Fragment derselben ist.

20 4. Verbindung mit der Formel:



in der

eine der Gruppen Z¹, Z², Z³ und Z⁴ RCW(HNV)_nY ist und die anderen H₂ oder =O sind;

35 R ein zweibindiger Rest mit 1 bis 6 Kohlenstoffatomen und 0 bis 2 Heteroatomen ist;

W = NH oder =O ist, mit der Bedingung, daß das an das mit Y verknüpfte Kohlenstoffatom gebundene W H₂ ist, das an das an Y gebundene Kohlenstoffatom gebunden ist, wenn Y -NH₂ ist;

40 V gleich oder verschieden von RCW ist und unter die Definitionen von RCW fällt;

n 0 oder 1 ist;

45 T eine entfernbare Schutzgruppe, Wasserstoff oder ein Alkalimetallion ist;

Y eine organische, einen aktiven Ester bildende Oxyverbindung, -NH₂, -NHNH₂ oder ein Polypeptid aus mindestens zwei Aminosäuren ist;

50 X eine Bindung, Methylen oder CHZ⁴ ist; und

die Gruppen A, die gleich oder verschieden sein können, Wasserstoff oder niederes Alkyl mit 1 bis 6 Kohlenstoffatomen sind.

55 5. Verbindung nach Anspruch 4, in der X eine Bindung ist.

6. Verbindung nach Anspruch 4 oder 5, in der Y eine aktivierte Phenoxygruppe ist.

EP 0 188 256 B1

7. Verfahren zur Herstellung eines radioaktiv markierten Polypeptids mit den Merkmalen:

Zusammengeben eines Polypeptids und einer Verbindung mit der Formel gemäß Anspruch 1 in einem wäßrigen Medium, wobei Y die austretende Gruppe eines aktiven Esters, der eine Amid- oder Amidin-Verknüpfung mit dem Polypeptid bilden kann, oder NH_2 oder NHNH_2 ist, mit der Bedingung, daß, wenn Y NHNH_2 oder $-\text{NH}_2$ ist, das Polypeptid eine Oxo-Gruppe als Ergebnis einer Glykolspaltung eines an das Polypeptid gebundenen Zuckers aufweist und das Zusammengeben unter reduktiven Aminierungsbedingungen erfolgt;

wobei die Verbindung eine stabile, kovalente Bindung mit dem Polypeptid unter weitgehender Anwesenheit von unchelatisiertem Radionuklid bildet.


8. Verfahren zur Herstellung eines radioaktiv markierten Polypeptids mit den Merkmalen:

Zusammengeben einer Verbindung gemäß einem der Ansprüche 4 bis 6, in der T ein Alkalimetallion oder eine Schwefel-Schutzgruppe und Y das Polypeptid ist, und eines Metall-Radionuklids in einer reduzierten, chelatisierten, veränderlichen Form in einem wäßrigen Medium;

wobei das Radionuklid durch die Verbindung chelatisiert wird; und


Waschen der das Radionuklid enthaltenden Verbindung, um unchelatisiertes, an die Verbindung gebundenes Radionuklid zu entfernen.

9. Verbindung mit der Formel



in der T eine Schwefel-Schutzgruppe ist.

10. Verbindung mit der Formel:




in der M ein Metall-Radionuklid, ausgewählt aus $^{99\text{m}}\text{Tc}$ und ^{186}Re , darstellt.

Patentansprüche für folgenden Vertragsstaat: AT

1. Verfahren zur Herstellung einer Verbindung mit der Formel:

EP 0 188 256 B1



in der

- 15 M ein chelatisierbares Metall- oder Metalloxyd-Radionuklidion ist;
eine der Gruppen Z^1 , Z^2 , Z^3 und Z^4 $R'CW'(HNV)_nY'$ ist und die anderen H_2 oder $=O$ sind;
 R' ein aliphatischer, zweibindiger Rest mit 1 bis 6 Kohlenstoffatomen und 0 bis 2 Heteroatomen ist;
- 20 Y' die austretende Gruppe eines aktiven Esters, $-NH_2$, $-NHNH_2$ oder ein Polypeptid mit einem Molekulargewicht von mindestens ungefähr 1000 ist;
 V' die Formel $R'CW'$ aufweist;
- 25 $W' = NH$ oder $=O$ ist, mit der Bedingung, daß das an das mit Y' verknüpfte Kohlenstoffatom gebundene W' H_2 ist, das an das an Y' gebundene Kohlenstoffatom gebunden ist, wenn $Y' = -NH_2$ ist;
 X' eine Bindung, Methylen oder CHZ^4 ist;
- 30 die Gruppen A, die gleich oder verschieden sein können, Wasserstoff oder niederes Alkyl mit 1 bis 6 Kohlenstoffatomen sind; und
 n' 0 oder 1 ist,
- 35 mit dem Schritt des Chelatisierens eines Metall- oder Metalloxyd-Radionuklid-Ions mit einem Vorläufer der Verbindung.
2. Verfahren nach Anspruch 1, in dem Y' ein Polypeptid ist.
- 40 3. Verfahren nach Anspruch 2, dadurch gekennzeichnet, daß Y' ein Immunoglobulin oder spezifisches bindendes Fragment desselben ist.
- 45 4. Verfahren zur Herstellung eines radioaktiv markierten Polypeptids mit den Merkmalen:
Zusammengeben eines Polypeptids und einer Verbindung mit der Formel gemäß Anspruch 1 in einem wäßrigen Medium, wobei Y die austretende Gruppe eines aktiven Esters, der eine Amid- oder Amidin-Verknüpfung mit dem Polypeptid bilden kann, oder NH_2 oder $NHNH_2$ ist, mit der Bedingung, daß, wenn Y $NHNH_2$ oder $-NH_2$ ist, das Polypeptid eine Oxo-Gruppe als Ergebnis einer Glykolspaltung eines an das Polypeptid gebundenen Zuckers aufweist und das Zusammengeben unter reduktiven Aminierungsbedingungen erfolgt;
- 50 wobei die Verbindung eine stabile, kovalente Bindung mit dem Polypeptid unter weitgehender Abwesenheit von unchelatisiertem Radionuklid bildet.
- 55 5. Verfahren zur Herstellung eines radioaktiv markierten Polypeptids mit dem Merkmal:
Waschen einer nach Anspruch 1 erhaltenen, ein Radionuklid enthaltenden Verbindung, um unchelati-


EP 0 188 256 B1

siertes, an die Verbindung gebundenes Radionuklid zu entfernen.

6. Verfahren nach einem jeden der vorherigen Ansprüche, dadurch gekennzeichnet, daß eine Verbindung mit der Formel:

5

10



15

hergestellt wird,

in der M ein Metall-Radionuklid, ausgewählt aus ^{99m}Tc und ¹⁸⁶Re, darstellt.

20

25

30

35

40

45

50

55